## **Product Information**



# bertinHistone H3 Dimethyl (K)-Phospho (S) Monoclonal Antibody

Cat No: G01056 - 100 µL

General Data	and the second
Shipping:	wet ice
Formulation:	Liquid Ascite, does not contain any preservative therefore avoid repeat freezing and thawing cycles
Host:	Mouse
Antigen:	Ovalbumin conjugated synthetic peptide ASGSSRSKS[dimethyl K]DRKYTL(C) corresponding to aa 180-195 of human TAF10. The generated 51TA2H12 clone recognizes S10- or S28-phosphorylated histone H3, but only when doubly modified with H3K9me2 or H3K27me2 (see re
Clone:	2H12
Isotype:	lgG1, κ
Application(s):	ELISA
	Western Blot
Application(s):	Immunofluorescence
	Immunoprecipitation
	Recommended dilutions: 1/500-1/5000
Specificity:	dimethylated (K9)-phosphorylated (S10) histone H3 or dimethylated (K27)- phosphorylate (S28) histone H3. These histone H3 modifications are correlated with mitosis.
<b>.</b>	

Scientific Literature

Eberlin A, Grauffel C, Oulad-Abdelghani M, Robert F, Torres-Padilla ME, Lambrot R, Spehner D, Ponce-Perez L, Würtz JM, Stote RH, Kimmins S, Schultz P, Dejaegere A, Tora L. Histone H3 tailscontaining dimethylated lysine and adjacent phosphorylated serine modifications adopt a specific conformation during mitosis and meiosis. Mol Cell Biol. 2008 Mar;28(5):1739-54.

#### FP/16/24

#### For research laboratory use only â€" Not for human diagnostic use.

Buyers agree to purchase the material subject to Purchasing Terms that can be found on our website. Seek appropriate training to safely handle this product under normal conditions. Use

Bertin Bioreagent does not make any other warranty or representation whatsoever whether expressed or implied, with respect to these products. In no event will Bertin Bioreagent be liable

### Contact Bertin Bioreagent

Parc d'Activités du Pas du Lac 10 bis avenue AmpÃ<sup>-</sup>re 78180 Montigny le Bretonneux - France Tel.: +33 (0)139 306 036 https://www.bertin-bioreagent.com/pa206/contact-us https://www.bertin-bioreagent.com