Product Information





General Data

Shipping: wet ice

Formulation: Liquid Ascite, does not contain any preservative therefore

avoid repeat freezing and thawing cycles

Host: Mouse

Antigen: Synthetic peptide

Clone: 1B5

Isotype: $IgG1, \kappa$

Application(s): ELISA

Western Blot

Immunoprecipitation

Immunocytochemistry

Recommended dilutions: 1/500-1/5000

Specificity: human XPG.

Product Overview

ERCC5 or XPG is a single-strand specific DNA endonuclease that makes the 3' incision in DNA excision repair following UV-induced damage. The protein may also function in other cellular processes, including RNA polymerase II transcription, and transcription-coupled DNA repair. Mutations cause xeroderma pigmentosum complementation group G (XP-G), which is also referred to as xeroderma pigmentosum VII (XP7), a skin disorder characterized by hypersensitivity to UV light and increased susceptibility for skin cancer development following UV exposure. Some patients also develop Cockayne syndrome.

Scientific Literature

Riedl T, Hanaoka F, Egly JM. The comings and goings of nucleotide excision repair factors on damaged DNA. EMBO J. 2003 Oct 1;22(19):5293-303.

FP/18/24

For research laboratory use only â€" Not for human diagnostic use.

Buyers agree to purchase the material subject to Purchasing Terms that can be found on our website. Seek appropriate training to safely handle this product under normal conditions. Use the recommended personal protective equipment to prevent chemical exposures.

Bertin Bioreagent does not make any other warranty or representation whatsoever whether expressed or implied, with respect to these products. In no event will Bertin Bioreagent be liable for incidental, consequential or punitive damages.

Contact Bertin Bioreagent

Parc d'Activités du Pas du Lac
10 bis avenue AmpÃ"re
78180 Montigny le Bretonneux - France
Tel.: +33 (0)139 306 036
https://www.bertin-bioreagent.com/pa206/contact-us
https://www.bertin-bioreagent.com