



Precellys Multi Tissue Dissociation Kit (mouse)



USER GUIDE

Precellys Multi Tissue Dissociation Kit (mouse)

For research laboratory use only



DISCLAIMER

Bertin Technologies and/or its subsidiaries accept no responsibility with respect to the present document, whether explicit or implicit, in particular, but not limited to commercial warranties or individual warranties.

In the absence of legislation to the contrary, neither Bertin Technologies nor its subsidiaries shall be held liable, either in contract or in tort, in terms of warranty or any provision, for any damage arising in particular but not exclusively during the use of this document.

Neither Bertin Technologies nor its employees shall be held responsible for losses or costs arising from said use.

The information contained in the present document is intended solely for use with this Bertin Technologies product. Bertin Technologies shall not be liable for any use of this information in its application to other products.

Bertin Technologies reserves the right to revise the present document and make periodic modifications without prior warning.

This liability disclaimer clause is governed by and interpreted in accordance with French law. Any dispute arising from the content of said clause is therefore subject to the exclusive jurisdiction of the Courts of Paris, France.

PROPRIETARY RIGHTS AND CONFIDENTIALITY

The information, research, drawings, and diagrams contained in the present document remain the property of Bertin Technologies and are confidential.

The information contained in the present document may not be used (whether in part or in full), disclosed, or reproduced without prior authorization from Bertin Technologies.

TABLE OF CONTENTS

1.	PRECAUTION FOR USE
2.	MATERIALS AND EQUIPMENT REQUIRED
3.	REAGENT PREPARATION
4.	DISSOCIATION OF MOUSE LIVER
5.	DISSOCIATION OF MOUSE LUNG
6.	DISSOCIATION OF MOUSE HEART
7.	DISSOCIATION OF MOUSE SPLEEN

BERTIN TECHNOLOGIES
Parc d'activités du Pas du Lac
10 bis, avenue Ampère
78180 Montigny le Bretonneux
France
Tél: + 33 (0)1 39 30 61 60

Precellys Multi Tissue Dissociation Kit (mouse) is a registered trade mark of Bertin Technologies.

© 2024 Bertin Technologies. All rights reserved

Document reference: 24700-800-DU001-C

Expiry date: stated on the package

This kit contains:

Designation	Item #	Contence	Quantity
Precellys Dissociation tubes	D30717	Tubes	20
Precellys Multi-Tissue Dissociation Kit	D05717	Enzyme MT-7	4x square bottles
		Enzyme MT-8	2x 2,5 mL Tubes

The package contains enough for 20 reactions.

1. PRECAUTION FOR USE

Temperature

- ▶ Precellys Dissociation tubes must be stored at room temperature.
- ▶ Before opening the Precellys Multi-Tissue Dissociation Kit must be stored -20°C
- ▶ Enzyme MT7/Enzyme MT8 must be aliquots of appropriate volume to avoid thaw-cycles and store the aliquots at -20°C during maximum 6 months.

2. MATERIALS AND EQUIPMENT REQUIRED

In addition to standard laboratory equipment, the following materials may be required:

- ▶ Precision micropipettes (20 to 1000 µL)
- ▶ Incubator
- ▶ Precellys Evolution (P002511-PEVT0-A.0)
- ▶ 15 mL tube holder on the Precellys Evolution (S000915-PEVO0-A)
- ▶ DMEM/F-12, HEPES
- ▶ Fetal Bovine Serum (FBS)
- ▶ Phosphate-Buffered Saline (PBS) pH 7.4
- ▶ Red Blood Cell (RBC) Lysis buffer 1X
- ▶ PEB buffer: Prepare a solution containing phosphate-buffered saline (PBS), pH 7.2, 0.5% Bovine Serum Albumin (BSA), and 2 mM EDTA.
- ▶ Debris Removal Solution (optional).

In function of the type of the tissue, the material and equipment required is not the same. For more information go to the specific tissue protocol.

In this protocol, some steps will ask to run the following program on the Precellys Evolution instrument: 4500 rpm x 3 seconds.

Please note that if you don't have the latest software version you will have to manually stop the run at 3 seconds, as the minimum run in this version is 5 seconds.

Please refer to your local distributor to upgrade the software.

3. REAGENT PREPARATION

The dissociation enzymes (MT7/MT8) are ready to use solution.

**4. DISSOCIATION OF MOUSE LIVER**

1. Prepare enzyme mix for dissociation by pipetting the following reagents into a Precellys Dissociation Tube: 4875 µL of enzyme MT-7 and 125 µL of enzyme MT-8.
2. Transfer liver sample into the Precellys Dissociation Tube containing the enzyme mix. Close tightly.
3. Place the Precellys Dissociation Tube in the appropriate 15 mL tube holder on the Precellys Evolution.
4. Run the following program: 4500 rpm x 3 seconds.
5. Remove tube from Precellys Evolution and incubate sample for 30 minutes at 37 °C.
6. After incubation, place back the sample in the Precellys Evolution. Run the following program: 4500 rpm x 3 seconds.
7. Recover Precellys Dissociation Tube.
8. Resuspend sample and filter through a 100 µm cell strainer placed over a 50 mL conical tube.
9. Wash the filter with 5 mL of DMEM/F-12, HEPES.
10. Discard the filter and centrifuge cell suspension at 300xg for 10 minutes. Aspirate supernatant completely.
11. Resuspend cells with an appropriate buffer of the required volume for downstream application.

5. DISSOCIATION OF MOUSE LUNG

1. Prepare enzyme mix for dissociation by pipetting the following reagents into a Precellys Dissociation Tube: 4875 µL of enzyme MT-7 and 125 µL of enzyme MT-8.
2. Transfer lung sample into the Precellys Dissociation Tube containing the enzyme mix. Close tightly.
3. Place the Precellys Dissociation Tube in the appropriate 15 mL tube holder on the Precellys Evolution.
4. Run the following program: 4500 rpm x 3 seconds.
5. Remove tube from Precellys Evolution and incubate sample for 30 minutes at 37 °C.
6. After incubation, place back the sample in the Precellys Evolution. Run the following program: 4500 rpm x 3 seconds.
7. Recover Precellys Dissociation Tube.
8. Resuspend sample with 7.5 mL of DMEM/F-12, HEPES + 20% FBS.
9. Filter through a 70 µm cell strainer placed over a 50 mL conical tube.
10. Wash the filter with 3 mL of DMEM/F-12, HEPES + 20% FBS.
11. Discard the filter and centrifuge cell suspension at 600xg for 5 minutes. Aspirate supernatant completely.
12. Resuspend cells with an appropriate buffer to the required volume for downstream applications



6. DISSOCIATION OF MOUSE HEART

1. Prepare enzyme mix for dissociation by pipetting the following reagents into a Precellys Dissociation Tube: 4875 µL of enzyme MT-7 and 125 µL of enzyme MT-8.
2. Transfer heart sample into the Precellys Dissociation Tube containing the enzyme mix. Close tightly.
3. Place the Precellys Dissociation Tube in the appropriate 15 mL tube holder on the Precellys Evolution.
4. Run the following program: 4500 rpm x 3 seconds.
5. Remove tube from Precellys Evolution and incubate sample for 30 minutes at 37 °C.
6. After incubation, place back the sample in the Precellys Evolution. Run the following program: 4500 rpm x 5 seconds.
7. Recover Precellys Dissociation Tube.
8. Resuspend sample with 7.5 mL of DMEM/F-12, HEPES + 20% FBS.
9. Filter through a 70 µm cell strainer placed over a 50 mL conical tube.
10. Wash the filter with 3 mL of DMEM/F-12, HEPES + 20% FBS.
11. Discard the filter and centrifuge cell suspension at 600xg for 5 minutes. Aspirate supernatant completely.
12. Proceed with Debris removal step (optional - not included).
13. Proceed with Red blood cell lysis (section 6.2.3)

► Red blood cell lysis

1. Resuspend cell pellet in 1 mL of PEB buffer and add 10 mL of 1X Red Blood Cell Lysis buffer.
2. Incubate for 2 minutes at room temperature.
3. Centrifuge at 600xg for 5 minutes. Aspirate supernatant completely.
4. Resuspend cell pellet in 10 mL of PBS.
5. Centrifuge at 600xg for 5 minutes. Aspirate the supernatant completely.
6. Resuspend cells with an appropriate buffer to the required volume for downstream applications.

7. DISSOCIATION OF MOUSE SPLEEN/THYMUS


1. Prepare enzyme mix for dissociation by pipetting the following reagents into a Precellys Dissociation Tube: 4875 µL of enzyme MT-7 and 125 µL of enzyme MT-8.
2. Transfer spleen sample into the Precellys Dissociation Tube containing the enzyme mix. Close tightly.
3. Recover the dissociation tubes from Precellys Evolution. Incubate sample for 15 minutes at 37 °C.
4. After incubation, place back the sample in the Precellys Evolution. Run the following program: 4500 rpm x 2 cycles of 5 seconds, with a 10 second pause between cycles.
5. Recover the Precellys Dissociation Tube.
6. Resuspend sample with 7.5 mL of DMEM/F-12, HEPES and filter through a 30 µm cell strainer placed over a 15 mL conical tube.
7. Wash the filter with 2.5 mL of DMEM/F-12, HEPES.
8. Discard the filter and centrifuge cell suspension at 300xg for 10 minutes. Aspirate supernatant completely.
9. Resuspend cells with an appropriate buffer to the required volume for downstream applications.



**Do not hesitate to contact our
after-sales services for further information at**

**+33 (0)139 306 036
tech@bertin-bioreagent.com**

EU webstore: Bertin-bioreagent.com
US webstore: Bertin-corp.com



Bertin Technologies
10 avenue Ampère, 78180 Montigny-le-Bretonneux, France
www.bertin-technologies.com